Author Index

Akiyama, Y., see Takeda, N. 558(1991)175
Alderweireldt, F. C., see Maes, P. C. 558(1991)343
Alexander, P. W., see Buchberger, W. 558(1991)181
Algiman, E., see Kong Sing, Y. L. 558(1991)43
Alvi, S. N., see Husain, S. 558(1991)435
Araki, T., see Kimata, K. 558(1991)19
Baastoe, M. B. and Lundanes, E.
Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458
Baba, Y., Matsuura, T., Wakamoto, K. and Tsuhako, M.
Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273
Baillet, A., see Taverna, M. 558(1991)105
Baylocq-Ferrier, D., see Taverna, M. 558(1991)105
Begutter, H., see Pachinger, A. 558(1991)369
Benlyas, M., see Moussaoui, M. 558(1991)71
Beran, M., Přikrylová, V., Sedmera, P., Novák, J., Zima, J., Blumauerová, M. and Todorov, T. K.
Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265
Bertrand, O., see Kong Sing, Y. L. 558(1991)43
Bhatti, A. G., see Khuhawar, M. Y. 558(1991)187
Black, R. M., Clarke, R. J. and Read, R. W.
Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio) ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)405
Black, R. M. and Read, R. W.
Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)393
Blay, P., see Pleasance, S. 558(1991)155
Blumauerová, M., see Beran, M. 558(1991)265
Bosman, J., see Witte, D. T. 558(1991)333
Both, D. A. and Jemal, M.
Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257
Brilis, G. M., see Brumley, W. C. 558(1991)223
Brownrigg, C. M., see Brumley, W. C. 558(1991)223
Brumley, W. C., Brownrigg, C. M. and Brilis, G. M.
Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography–mass spectrometry after fractionation 558(1991)223
Buchberger, W., Haddad, P. R. and Alexander, P. W.
Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181
Burns, K. I., see Elchuk, S. 558(1991)197
Carelse, O., Chetsanga, C. J. and Mubumbila, M. V.
Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318
Caruel, H., Rigal, L. and Gaset, A.
Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar–cation complexes and the elution order 558(1991)89
Cassidy, R. M., see Elchuk, S. 558(1991)197
Cebrian Perez, J. A., see Ortin, A. 558(1991)357
Chang, F. C., see Iida, T. 558(1991)451
Chen, S., Ocampo, A. P. and Kucera, P. J.
Liquid chromatographic method for the determination of calcium cyanamide using pre-column derivatization 558(1991)141
Chetsanga, C. J., see Carelse, O. 558(1991)318
Chiari, M., Giacomini, M., Micheletti, C. and Righetti, P. G.
Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285
Clarke, R. J., see Black, R. M. 558(1991)405
Cremer, J., Meyer, H. and Henning, K.
Separation of methylated and non-methylated cyanoginosin-LR homologues of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 by reversed-phase medium-pressure liquid chromatography 558(1991)430
De Boer, J. H., see Witte, D. T. 558(1991)333
De Graaf, G., see Dos Ramos, F. J. 558(1991)125
Dewaele, C., see Maes, P. C. 558(1991)343
De Zeeuw, R. A., see Witte, D. T. 558(1991)333
Dhermy, D., see Kong Sing, Y. L. 558(1991)43

Author Index

Akiyama, Y., see Takeda, N. 558(1991)175
Alderweireldt, F. C., see Maes, P. C. 558(1991)343
Alexander, P. W., see Buchberger, W. 558(1991)181
Algiman, E., see Kong Sing, Y. L. 558(1991)43
Alvi, S. N., see Husain, S. 558(1991)435
Araki, T., see Kimata, K. 558(1991)19
Baastoe, M. B. and Lundanes, E.
Capillary supercritical fluid chromatography of primary aliphatic amines using carbon dioxide and nitrous oxide as mobile phases 558(1991)458
Baba, Y., Matsuura, T., Wakamoto, K. and Tsuhako, M.
Comparison of high-performance liquid chromatography with capillary gel electrophoresis in single-base resolution of polynucleotides 558(1991)273
Baillet, A., see Taverna, M. 558(1991)105
Baylocq-Ferrier, D., see Taverna, M. 558(1991)105
Begutter, H., see Pachinger, A. 558(1991)369
Benlyas, M., see Moussaoui, M. 558(1991)71
Beran, M., Přikrylová, V., Sedmera, P., Novák, J., Zima, J., Blumauerová, M. and Todorov, T. K.
Isolation of erythromycin A N-oxide and pseudoerythromycin A hemiketal from fermentation broth of *Saccharopolyspora erythraea* by thin-layer and high-performance liquid chromatography 558(1991)265
Bertrand, O., see Kong Sing, Y. L. 558(1991)43
Bhatti, A. G., see Khuhawar, M. Y. 558(1991)187
Black, R. M., Clarke, R. J. and Read, R. W.
Analysis of 1,1'-sulphonylbis[2-(methylsulphinyl)ethane] and 1-methylsulphinyl-2-[2-(methylthio) ethylsulphonyl]ethane, metabolites of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)405
Black, R. M. and Read, R. W.
Methods for the analysis of thiodiglycol sulphoxide, a metabolite of sulphur mustard, in urine using gas chromatography–mass spectrometry 558(1991)393
Blay, P., see Pleasance, S. 558(1991)155
Blumauerová, M., see Beran, M. 558(1991)265
Bosman, J., see Witte, D. T. 558(1991)333
Both, D. A. and Jemal, M.
Stereoisomeric purity determination of captopril by capillary gas chromatography 558(1991)257
Brilis, G. M., see Brumley, W. C. 558(1991)223
Brownrigg, C. M., see Brumley, W. C. 558(1991)223
Brumley, W. C., Brownrigg, C. M. and Brilis, G. M.
Characterization of nitrogen-containing aromatic compounds in soil and sediment by capillary gas chromatography–mass spectrometry after fractionation 558(1991)223
Buchberger, W., Haddad, P. R. and Alexander, P. W.
Separation of metal complexes of ethylenediaminetetraacetic acid in environmental water samples by ion chromatography with UV and potentiometric detection 558(1991)181
Burns, K. I., see Elchuk, S. 558(1991)197
Carelse, O., Chetsanga, C. J. and Mubumbila, M. V.
Purification of common bean chloroplast DNA by DEAE cellulose column chromatography 558(1991)318
Caruel, H., Rigal, L. and Gaset, A.
Carbohydrate separation by ligand-exchange liquid chromatography. Correlation between the formation of sugar–cation complexes and the elution order 558(1991)89
Cassidy, R. M., see Elchuk, S. 558(1991)197
Cebrian Perez, J. A., see Ortin, A. 558(1991)357
Chang, F. C., see Iida, T. 558(1991)451
Chen, S., Ocampo, A. P. and Kucera, P. J.
Liquid chromatographic method for the determination of calcium cyanamide using pre-column derivatization 558(1991)141
Chetsanga, C. J., see Carelse, O. 558(1991)318
Chiari, M., Giacomini, M., Micheletti, C. and Righetti, P. G.
Synthesis of a new acrylamido buffer (acryloylhistamine) for isoelectric focusing in immobilized pH gradients and its analysis by capillary zone electrophoresis 558(1991)285
Clarke, R. J., see Black, R. M. 558(1991)405
Cremer, J., Meyer, H. and Henning, K.
Separation of methylated and non-methylated cyanoginosin-LR homologues of the cyanobacterium *Microcystis aeruginosa* strain PCC 7806 by reversed-phase medium-pressure liquid chromatography 558(1991)430
De Boer, J. H., see Witte, D. T. 558(1991)333
De Graaf, G., see Dos Ramos, F. J. 558(1991)125
Dewaele, C., see Maes, P. C. 558(1991)343
De Zeeuw, R. A., see Witte, D. T. 558(1991)333
Dhermy, D., see Kong Sing, Y. L. 558(1991)43

Diallo, B., Vanhaelen-Fastré, R. and Vanhaelen, M.
Direct coupling of high-speed counter-current chromatography to thin-layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica* 558(1991)446
Doornbos, D. A., see Witte, D. T. 558(1991)333
Dos Ramos, F. J., Noronha da Silveira, I. and De Graaf, G.
Column liquid chromatographic determination of carbadox and olaquindox in feeds 558(1991)125
Eisner, E., see Pachinger, A. 558(1991)369
Elchuk, S., Burns, K. I., Cassidy, R. M. and Lucy, C. A.
Reversed-phase separation of transition metals, lanthanides and actinides by elution with mandelic acid 558(1991)197
Fadeev, A. Y., Lisichkin, G. V., Runov, V. K. and Staroverov, S. M.
Diffusion of sorbed pyrene in the bonded layer of reversed-phase silicas. Effect of alkyl chain length and pore diameter 558(1991)31
Fowler, W. K., Stewart, D. C., Weinberg, D. S. and Sarver, E. W.
Gas chromatographic determination of the lewisite hydrolysate, 2-chlorovinylarsonous acid, after derivatization with 1,2-ethanedithiol 558(1991)235
Gardana, C., see Pietta, P. 558(1991)296
Gaset, A., see Caruel, H. 558(1991)89
Gasparič, J. and Skutil, J.
Comparative study of chromatography on thin layers impregnated with organic stationary phases. Chromatographic separation of nitrophenols 558(1991)415
Gebauer, P. and Thormann, W.
Isotachophoresis of proteins in uncoated open-tubular fused-silica capillaries with a simple approach for column conditioning 558(1991)423
Giacomini, M., see Chiari, M. 558(1991)285
Goto, J., see Iida, T. 558(1991)451
Gutiérrez, M. C., see Linares, P. 558(1991)147
Haddad, P. R., see Buchberger, W. 558(1991)181
Haginaka, J., see Kimata, K. 558(1991)19
Hahn, A., see Sander, S. 558(1991)115
Henning, K., see Cremer, J. 558(1991)430
Highsmith, F. A., see Orthner, C. L. 558(1991)55
Hosoya, K., see Kimata, K. 558(1991)19
Husain, S., Nageswara Sarma, P., Swamy, N. S., Alvi, S. N. and Nageswara Rao, R.
High-performance liquid chromatographic separation and determination of small amounts of 2-naphthaleneacetic acid in 1-naphthaleneacetic acid 558(1991)435
Iida, T., Tamaru, T., Chang, F. C., Goto, J. and Nambara, T.
Capillary gas chromatographic analysis of oxo, oxo-hydroxy and unsaturated bile acid glycine conjugates 558(1991)451
Jegorov, A., Trnka, T. and Stuchlík, J.
High-performance liquid chromatographic detection of enantiomeric amino alcohols after derivatization with *o*-phthaldialdehyde and various thiosugars 558(1991)311
Jemal, M., see Both, D. A. 558(1991)257
Jönsson, B., Welinder, H. and Skarping, G.
Determination of hexahydrophthalic anhydride in air using gas chromatography 558(1991)247
Kavanagh, P. E., see Shalliker, R. A. 558(1991)440
Khuhawar, M. Y. and Bhatti, A. G.
Thermoanalytical and chromatographic studies of copper(II), nickel(II) and oxovanadium(IV) complexes of tetradentate Schiff bases derived from *β*-diketones and 2,3-diaminopentane 558(1991)187
Kimata, K., Hosoya, K., Tanaka, N., Araki, T., Tsuboi, R. and Haginaka, J.
Effect of stationary phase structure on retention and selectivity of restricted-access reversed-phase packing materials 558(1991)19
Klus, H., see Pachinger, A. 558(1991)369
Kohata, K., Watanabe, M. and Yamanaka, K.
Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography 558(1991)131
Kong Sing, Y. L., Algiman, E., Kroviarski, Y., Massot, C., Dhermy, D. and Bertrand, O.
Modification of reversed-phase columns with dyed surfactants. Preparation of mechanically resistant efficient immobilized dyes for protein purification 558(1991)43
Kroviarski, Y., see Kong Sing, Y. L. 558(1991)43
Kucera, P. J., see Chen, S. 558(1991)141
Lanin, S. N. and Nikitin, Y. S.
Retention data for five ketotrichothecenes in reversed-phase high-performance liquid chromatography with different eluent systems 558(1991)81
Lázaro, F., see Linares, P. 558(1991)147
Lepojević, Ž., see Vrbaški, L. 558(1991)328
Linares, P., Gutiérrez, M. C., Lázaro, F., Luque de Castro, M. D. and Valcárcel, M.
Determination of benzocaine, dextromethorphan and cetylpyridinium ion by high-performance liquid chromatography with UV detection 558(1991)147
Lisichkin, G. V., see Fadeev, A. Y. 558(1991)31
López-Pérez, M. J., see Ortin, A. 558(1991)357
Lucy, C. A., see Elchuk, S. 558(1991)197
Lundanes, E., see Baastoe, M. B. 558(1991)458

Luque de Castro, M. D., see Linares, P. 558(1991)147
Madurawe, R. D., see Orthner, C. L. 558(1991)55
Maes, P. C., Nagels, L. J., Dewaele, C. and Alderweireldt, F. C.
Characterization of cellulase-based enzyme reactors for the high-performance liquid chromatographic determination of β-D-glucan oligosaccharides 558(1991)343
Massot, C., see Kong Sing, Y. L. 558(1991)43
Matsuura, T., see Baba, Y. 558(1991)273
Mauri, P., see Pietta, P. 558(1991)296
Meyer, H., see Cremer, J. 558(1991)430
Micheletti, C., see Chiari, M. 558(1991)285
Morcol, T., see Orthner, C. L. 558(1991)55
Moussaoui, M., Benlyas, M. and Wahl, P.
Diffusion of proteins in the chromatographic gel AcA-34 558(1991)71
Mubumbila, M. V., see Carelse, O. 558(1991)318
Muiño Blanco, M. T., see Ortin, A. 558(1991)357
Nagels, L. J., see Maes, P. C. 558(1991)343
Nageswara Rao, R., see Husain, S. 558(1991)435
Nageswara Sarma, P., see Husain, S. 558(1991)435
Nambara, T., see Iida, T. 558(1991)451
Nikitin, D. I., see Řezanka, T. 558(1991)215
Nikitin, Y. S., see Lanin, S. N. 558(1991)81
Noronha da Silveira, I., see Dos Ramos, F. J. 558(1991)125
Novák, J., see Beran, M. 558(1991)265
Obrezkov, O. N., Shpigun, O. A., Zolotov, Y. A. and Shlyamin, V. I.
Ion chromatographic determination of anions, especially sulphur-containing anions, with conductimetric and kinetic detection 558(1991)209
Ocampo, A. P., see Chen, S. 558(1991)141
Oe, T., see Shimada, K. 558(1991)306
O'Hara, G., see Pleasance, S. 558(1991)155
Orthner, C. L., Highsmith, F. A., Tharakan, J., Madurawe, R. D., Morcol, T. and Velander, W. H.
Comparison of the performance of immunosorbents prepared by site-directed or random coupling of monoclonal antibodies 558(1991)55
Ortin, A., López-Pérez, M. J., Muiño Blanco, M. T. and Cebrian Perez, J. A.
Extraction of xanthine oxidase from milk by counter-current distribution in an aqueous two-phase system 558(1991)357
Pachinger, A., Eisner, E., Begutter, H. and Klus, H.
Determination of a metabolite of the herbicide pyridate in drinking and groundwater using high-performance liquid chromatography with amperometric detection 558(1991)369
Pietta, P., Gardana, C., Mauri, P. and Zecca, L.
High-performance liquid chromatographic analysis of flavonol glycosides of *Solidago virgaurea* 558(1991)296
Pirkle, W. H.
Unusual effect of temperature on the retention of enantiomers on a chiral column 558(1991)1
Pleasance, S., Blay, P., Quilliam, M. A. and O'Hara, G.
Determination of sulfonamides by liquid chromatography, ultraviolet diode array detection and ion-spray tandem mass spectrometry with application to cultured salmon flesh 558(1991)155
Přikrylová, V., see Beran, M. 558(1991)265
Quilliam, M. A., see Pleasance, S. 558(1991)155
Read, R. W., see Black, R. M. 558(1991)393
Read, R. W., see Black, R. M. 558(1991)405
Rehner, G., see Sander, S. 558(1991)115
Řezanka, T., Zlatkin, I. V., Viden, I., Slabova, O. I. and Nikitin, D. I.
Capillary gas chromatography–mass spectrometry of unusual and very long-chain fatty acids from soil oligotrophic bacteria 558(1991)215
Rigal, L., see Caruel, H. 558(1991)89
Righetti, P. G., see Chiari, M. 558(1991)285
Runov, V. K., see Fadeev, A. Y. 558(1991)31
Russell, I. M., see Shalliker, R. A. 558(1991)440
Sander, S., Hahn, A., Stein, J. and Rehner, G.
Comparative studies on the high-performance liquid chromatographic determination of thiamine and its phosphate esters with chloroethylthiamine as an internal standard using pre- and post-column derivatization procedures 558(1991)115
Sarver, E. W., see Fowler, W. K. 558(1991)235
Sedmera, P., see Beran, M. 558(1991)265
Shalliker, R. A., Kavanagh, P. E. and Russell, I. M.
Reversed-phase separations of high-molecular-weight polystyrenes in a dichloromethane–acetonitrile solvent system 558(1991)440
Shimada, K., Oe, T. and Suzuki, M.
Effect of derivatization of steroids on their retention behaviour in inclusion chromatography using cyclodextrin as a mobile phase additive 558(1991)306
Shlyamin, V. I., see Obrezkov, O. N. 558(1991)209
Shpigun, O. A., see Obrezkov, O. N. 558(1991)209
Skarping, G., see Jönsson, B. 558(1991)247
Skutil, J., see Gasparič, J. 558(1991)415
Slabova, O. I., see Řezanka, T. 558(1991)215
Ślebioda, M., see Wodecki, Z. J. 558(1991)302

Smith, R. M. and Wang, R.
Retention prediction of analytes in reversed-phase high-performance liquid chromatography based on molecular structure. VII. Separations using tetrahydrofuran–buffer eluents 558(1991)7
Staroverov, S. M., see Fadeev, A. Y. 558(1991)31
Stein, J., see Sander, S. 558(1991)115
Stewart, D. C., see Fowler, W. K. 558(1991)235
Storr-Hansen, E.
Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration 558(1991)375
Stuchlik, J., see Jegorov, A. 558(1991)311
Sudoh, Y., see Yakabe, Y. 558(1991)323
Suzuki, M., see Shimada, K. 558(1991)306
Swamy, N. S., see Husain, S. 558(1991)435
Takahata, Y., see Yakabe, Y. 558(1991)323
Takeda, N. and Akiyama, Y.
Pre-column derivatization of sulfa drugs with fluorescamine and high-performance liquid chromatographic determination at their residual levels in meat and meat products 558(1991)175
Tamaru, T., see Iida, T. 558(1991)451
Tanaka, N., see Kimata, K. 558(1991)19
Taverna, M., Baillet, A., Werner, R. and Baylocq-Ferrier, D.
Liquid chromatographic method for the determination of the carbohydrate moiety of glycoproteins. Application to α_1-acid glycoprotein and tissue plasminogen activator 558(1991)105
Tharakan, J., see Orthner, C. L. 558(1991)55
Thormann, W., see Gebauer, P. 558(1991)423
Todorov, T. K., see Beran, M. 558(1991)265
Torlop, B., see Wodecki, Z. J. 558(1991)302
Trnka, T., see Jegorov, A. 558(1991)311
Tsuboi, R., see Kimata, K. 558(1991)19
Tsuhako, M., see Baba, Y. 558(1991)273
Valcárcel, M., see Linares, P. 558(1991)147
Vanhaelen, M., see Diallo, B. 558(1991)446
Vanhaelen-Fastré, R., see Diallo, B. 558(1991)446
Víden, I., see Řezanka, T. 558(1991)215
Velander, W. H., see Orthner, C. L. 558(1991)55
Vrbaški, L. and Lepojević, Ž.
Rapid identification of microbial starch degradation products from a complex nutrient medium by a thin-layer chromatographic method 558(1991)328
Wahl, P., see Moussaoui, M. 558(1991)71
Wakamoto, K., see Baba, Y. 558(1991)273
Wang, R., see Smith, R. M. 558(1991)7
Watanabe, M., see Kohata, K. 558(1991)131
Weinberg, D. S., see Fowler, W. K. 558(1991)235
Welinder, H., see Jönsson, B. 558(1991)247
Werner, R., see Taverna, M. 558(1991)105
Witte, D. T., Bosman, J., De Zeeuw, R. A., De Boer, J. H. and Doornbos, D. A.
Optimization of the direct enantiomeric separation by high-performance liquid chromatography of the D-2 dopamine agonist N-0437 using response surface methodology and the multi-criteria decision-making method 558(1991)333
Wodecki, Z. J., Torlop, B. and Ślebioda, M.
Chromatographic determination of citric acid for monitoring the mould process 558(1991)302
Yakabe, Y., Sudoh, Y. and Takahata, Y.
Immobilization method for polyethylene glycol using a cross-linking co-agent 558(1991)323
Yamanaka, K., see Kohata, K. 558(1991)131
Zecca, L., see Pietta, P. 558(1991)296
Zima, J., see Beran, M. 558(1991)265
Zlatkin, I. V., see Řezanka, T. 558(1991)215
Zolotov, Y. A., see Obrezkov, O. N. 558(1991)209

Errata

J. Chromatogr., 539 (1991) 501–505.
Page 505: the authors would like to add the following "Acknowledgements" section: "The success of the work was critically dependent on an expression system which we received from Drs. Curtis L. Patton, Charles E. Egwuagu and James F. Young of the Yale University School of Medicine, New Haven, CT, USA."

J. Chromatogr., 554 (1991) 61–71.
Page 63, *Mobile phases*, line 4, "acetonitrile (ACN) (39%)" should read "acetonitrile (ACN) (30%)".